



Svetlana
ROMAN

GREEN'S FUNCTIONS FOR BOUNDARY-VALUE PROBLEMS WITH NONLOCAL BOUNDARY CONDITIONS

SUMMARY OF DOCTORAL DISSERTATION

PHYSICAL SCIENCES, MATHEMATICS (01 P)

VILNIUS 2011

VILNIUS UNIVERSITY

Svetlana Roman

GREEN'S FUNCTIONS FOR BOUNDARY-VALUE PROBLEMS
WITH NONLOCAL BOUNDARY CONDITIONS

Summary of Doctoral Dissertation
Physical Sciences, Mathematics (01 P)

Vilnius, 2011

The dissertation was prepared at Institute of Mathematics and Informatics, Vilnius University in 2007–2011.

Scientific supervisor

Prof Dr Artūras Štikonas (Vilnius University, Physical Sciences, Mathematics – 01 P).

The dissertation will be defended at the Council of Scientific Field of Mathematics at the Institute of Mathematics and Informatics of Vilnius University:

Chairman

Prof Dr Habil Mifodijus Sapagovas (Vilnius University, Physical Sciences, Mathematics – 01 P).

Members:

Prof Dr Habil Raimondas Čiegeis (Vilnius Gediminas Technical University, Physical Sciences, Mathematics – 01 P),

Prof Dr Habil Feliksas Ivanauskas (Vilnius University, Physical Sciences, Mathematics – 01 P),

Doc Dr Sigita Pečiulytė (Vytautas Magnus University, Physical Sciences, Mathematics – 01 P),

Prof Dr Habil Konstantinas Pileckas (Vilnius University, Physical Sciences, Mathematics – 01 P).

Opponents:

Prof Dr Vytautas Kleiza (Kaunas University of Technology, Physical Sciences, Mathematics – 01 P),

Prof Dr Stasys Rutkauskas (Vilnius University, Physical Sciences, Mathematics – 01 P).

The dissertation will be defended at the public meeting of the Council of Scientific of Mathematics in the auditorium number 203 at the Institute of Mathematics and Informatics of Vilnius University, at 1 p. m. on 14 December 2011.

Address: Akademijos st. 4, LT-08663 Vilnius, Lithuania.

The summary of the doctoral dissertation was distributed on 14th of November 2011.

A copy of the doctoral dissertation is available for review at the Library of Vilnius University.

VILNIAUS UNIVERSITETAS

Svetlana Roman

GRYNO FUNKCIJOS UŽDAVINIAMS SU NELOKALIOSIOMIS KRAŠTINĖMIS
SĄLYGOMIS

Daktaro disertacijos santrauka
Fiziniai mokslai, matematika (01 P)

Vilnius, 2011

Disertacija rengta 2007–2011 metais Vilniaus universiteto Mathematikos ir informatikos instituto.

Mokslinis vadovas

prof. dr. Artūras Štikonas (Vilniaus universitetas, fiziniai mokslai, matematika – 01 P).

**Disertacija ginama Vilniaus universiteto Matematikos ir informatikos instituto
Matematikos mokslo krypties taryboje:**

Pirmininkas

prof. habil. dr. Mifodijus Sapagovas (Vilniaus universitetas, fiziniai mokslai, matematika – 01 P).

Nariai:

prof. habil. dr. Raimondas Čiegeis (Vilniaus Gedimino technikos universitetas, fiziniai mokslai, matematika – 01 P),

prof. habil. dr. Feliksas Ivanauskas (Vilniaus universitetas, fiziniai mokslai, matematika – 01 P),

doc. dr. Sigita Pečiulytė (Vytauto Didžiojo universitetas, fiziniai mokslai, matematika – 01 P),

prof. habil. dr. Konstantinas Pileckas (Vilniaus universitetas, fiziniai mokslai, matematika – 01 P).

Oponentai:

prof. dr. Vytautas Kleiza (Kauno technologijos universitetas, fiziniai mokslai, matematika – 01 P),

prof. dr. Stasys Rutkauskas (Vilniaus universitetas, fiziniai mokslai, matematika – 01 P).

Disertacija bus ginama viešame Matematikos mokslo krypties tarybos posėdyje 2011 m. gruodžio mėn. 14 d. 13 val. Vilniaus universiteto Matematikos ir informatikos instituto 203 auditorijoje.

Adresas: Akademijos g. 4, LT-08663 Vilnius, Lietuva.

Disertacijos santrauka išsiuntinėta 2011 m. lapkričio mėn. 14 d.

Disertaciją galima peržiūrėti Vilniaus universiteto bibliotekoje.

Introduction

Problem formulation

In the dissertation Green's functions for the second-order and higher-order differential and difference equations with additional conditions, which are described by linearly independent linear functionals, are investigated. A lot of attention is given to the problems with nonlocal boundary conditions.

Topicality of the problem

In the theory of differential equations the basic concepts have been formulated studying the problems of classical mathematical physics. However, the modern problems motivate to formulate and investigate the new ones, for example, a class of nonlocal problems. Nonlocal conditions arise when we cannot measure data directly at the boundary. In this case, the problem is formulated where the value of the solution and/or a derivative is linked to a few points or the whole interval.

In 1963 J.R. Cannon [6] formulated new problem with boundary conditions which are now called *nonlocal* (N.I. Ionkin was the first who used the concept of *nonlocal boundary condition* in 1977 [24]). The similar problem was investigated by L.I. Kamynin in 1964 [27]. A.A. Samarskii and A.V. Bitsadze formulated the elliptical problem with nonlocal boundary conditions in 1969 [5]. Now some one-dimensional nonlocal conditions are called as Bitsadze–Samarskii conditions. Differential equations (for example, ordinary, elliptic, parabolic, etc.) with various types of nonlocal conditions were investigated by scientists: V.A. Il'in [16], E.I. Moiseev [17, 18], N.I. Ionkin [23, 25], A.V. Gulin [13, 12, 14, 15], J.R.L. Webb [51, 52], G. Infante [21, 22] and others. In Lithuania such problems were investigated by M. Sapagovas [10, 26, 31, 37, 38, 39, 40, 41, 42], R. Čiegeis [7, 8, 9], A. Štikonas [32, 33, 43, 45] and their students.

In this dissertation Green's functions of the second- and higher-order differential and difference equations with various conditions (for example, initial, boundary or nonlocal conditions) are investigated.

In mathematics, Green's function is a type of function used to solve nonhomogeneous equations subject to boundary conditions. Green's functions play an important role in

the theory of linear Ordinary and Partial Differential Equations. The term is used in physics, specifically in quantum field theory, electrodynamics and statistical field theory. It helps to investigate the existence and uniqueness of the solutions for many boundary problems.

Green's functions for problems with classical boundary conditions were analyzed by V.S. Vladimirov [48], U.M. Ascher, R.D. Russell and R.M.M. Mattheij [3], I. Stakgold [44], D.G. Duffy [11]. Besides A.A. Samarskii [34], E.S. Nikolaev [36], A.V. Gulin [35], N.S. Bahvalov, H.P. Židkov and G.M. Kobel'kov [4] studied Green's functions for problems with difference operator.

The investigation of the semi-linear problems with nonlocal boundary conditions and the existence of positive solutions are based on the investigation of Green's function for the linear problems with nonlocal boundary conditions.

Green's functions for the second- and higher-order boundary problems with various nonlocal conditions were constructed by D.R. Anderson [1, 2], J.R.L. Webb [21, 22, 49, 50], G. Infante [19, 20], R.Y. Ma [28, 29, 30], Y. Sun [46], L.X. Truong [47], B. Yang [54], Z. Zhao [55, 56] D. Xie, Y. Liu and C. Bai [53] and other scientists. In their works authors considered the existence and multiplicity of solutions applying various methods: lower and upper solution method, Leggett–Williams fixed-point theorem, Guo–Krasnoselskii fixed-point theorem, Leray–Schauder continuation principle, Avery–Peterson fixed-point theorem.

In the *first chapter* of this dissertation, the m -order linear differential equation with m additional conditions, which can be as classical as nonlocal, is considered. These conditions are formulated by means of independent linear functionals. Also in the first chapter, the formula, describing the relation between two problems with nonclassical and classical boundary conditions, is deduced.

In the *second chapter* the second-order differential equation is formulated. The results of the first chapter are applied to this problem. The examples show how Green's function for problems with varying complexity of nonlocal boundary conditions is constructed.

The differential equations, considered in the first and second chapters, can be approximated by the difference equations. In the *third chapter* the second-order difference equation with two conditions is investigated.

In the *fourth chapter* the results of the third chapter are generalized to the m -order difference equation with m additional conditions.

Research object

The main research object of the dissertation is differential and difference operators with various conditions, the solutions and Green's functions of these problems, the existence conditions of Green's functions and the application of the results for problems with nonlocal boundary conditions.

Aim and task of the work

The aim of this dissertation is to investigate the differential and discrete problems with additional conditions; to write down the expression of the solution for nonhomogeneous equation, if the fundamental solution is known; to find Green's function for these problems and to obtain a relation between two Green's functions for problems with the same equation and various conditions.

The main tasks of the work are:

1. to investigate linear space of the solutions for homogeneous equation;
2. to investigate the solutions of differential or discrete nonhomogeneous problems with the same equation, but with various conditions;
3. to find Green's functions for the differential and the difference problems and the existence conditions for them;
4. to investigate Green's functions for two problems with the same equation, but with various conditions;
5. to apply results to problems with nonlocal boundary conditions;
6. to compare results for differential or discrete problems.

Methodology of research

In the dissertation the method of variation of parameters is applied. Using this method and properties of determinants, Green's functions for differential and difference equations with variable coefficients and various boundary conditions have been obtained. Also we

describe linear functionals and functional determinants, which properties have been used. By means of packages “Maple” graphs are represented.

Scientific novelty

Green’s functions are often constructed to study the existence and multiplicity of the solutions for the second-order and higher-order semi-linear problems. In almost all cases Green’s function is searched for problem, which is a special case of the problem of this thesis.

In this thesis Green’s functions are investigated for problems with general linear conditions. The relation between two Green’s functions is obtained, which helps to investigate Green’s function for problem with nonlocal boundary conditions, if Green’s function for problem with classical boundary conditions is known. Mostly in the various monographs and articles the formulae of this type have been obtained for the problem with classical boundary conditions, but in this work they have been deduced for problem with any (linear) conditions. So, we can write the expression of Green’s function for problem with (for example, nonlocal) conditions if we know the fundamental basis of homogeneous equation and Green’s function of problem with other (for example, classical) conditions. The obtained results can be applied to problems with nonlocal boundary conditions.

Practical value

In the doctoral dissertation the obtained results might be applied for solving the differential or difference equations with the conditions of various (for example, initial, boundary, nonlocal conditions), and for investigating the existence, the uniqueness and the properties of the solutions.

Defended propositions

- The necessary and sufficient condition for the existence of Green’s function.
- The expressions of Green’s functions for the second- and higher-order differential and difference equations with various linear conditions.

- The relation between Green's function and Green's function for the problem with initial conditions.
- The relation between two Green's functions.
- The application of the obtained results to problems with nonlocal boundary conditions.

The scope of the scientific work

The doctoral dissertation consists of the introduction, four chapters, conclusions, the list of the references and the other one of author's publications. The total scope of the doctoral dissertation is 129 pages, 4 figures and 4 tables. The results of doctoral dissertation are published in 9 publications. The results were presented at 4 national and 5 international conferences. The language of the doctoral dissertation is Lithuanian.

Chapter 1. The m -Order Differential Problem

In the first chapter we investigate m -order differential equation with additional conditions

$$\mathcal{L}u := u^{(m)} + a^{m-1}(x)u^{(m-1)} + \cdots + a^1(x)u' + a^0(x)u = f(x), \quad (1)$$

$$\langle L_i, u \rangle = f_i \in \mathbb{K}, \quad i = 1, \dots, m, \quad (2)$$

where $a^i \in C[0, L]$, $i = 0, \dots, m - 1$, $f_i \in \mathbb{K} := \mathbb{R}, \mathbb{C}$, $i = 1, \dots, m$, $f \in C[0, L]$, and L_1, \dots, L_m are linearly independent functionals.

Let $\mathbf{L} = (L_1, \dots, L_m)$ and $\mathbf{w} = [w^1, \dots, w^m]$ be fundamental system of homogeneous equation (1), and we denote

$$H(x) := \begin{cases} 1, & x > 0, \\ 0, & x \leq 0, \end{cases}, \quad G^c(x, s) := H(x - s) \frac{\widetilde{W}[\mathbf{w}](x, s)}{W[\mathbf{w}](s)}, \quad x, s \in [0, L],$$

if $W[\mathbf{w}](s) \neq 0$, where

$$W[\mathbf{w}](s) := \begin{vmatrix} w^1(s) & (w^1)'(s) & \cdots & (w^1)^{(k-2)}(s) & (w^1)^{(k-1)}(s) \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ w^k(s) & (w^k)'(s) & \cdots & (w^k)^{(k-2)}(s) & (w^k)^{(k-1)}(s) \end{vmatrix}, \quad s \in [0, L],$$

$$\widetilde{W}[\mathbf{w}](x, s) := \begin{vmatrix} w^1(s) & (w^1)'(s) & \cdots & (w^1)^{(k-2)}(s) & w^1(x) \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ w^k(s) & (w^k)'(s) & \cdots & (w^k)^{(k-2)}(s) & w^k(x) \end{vmatrix}, \quad x, s \in [0, L].$$

We use functional $\langle \delta_x, w \rangle = w(x)$ and determinants:

$$D(\mathbf{L})[\mathbf{w}] := \begin{vmatrix} \langle L_1, w^1 \rangle & \cdots & \langle L_k, w^1 \rangle \\ \cdots & \cdots & \cdots \\ \langle L_1, w^k \rangle & \cdots & \langle L_k, w^k \rangle \end{vmatrix},$$

$$D(\mathbf{L}, \delta_x)[\mathbf{w}, G(\cdot, s)] := \begin{vmatrix} \langle L_1, w^1 \rangle & \cdots & \langle L_k, w^1 \rangle & w^1(x) \\ \cdots & \cdots & \cdots & \cdots \\ \langle L_1, w^k \rangle & \cdots & \langle L_k, w^k \rangle & w^k(x) \\ \langle L_1(\cdot), G(\cdot, s) \rangle & \cdots & \langle L_k(\cdot), G(\cdot, s) \rangle & G(x, s) \end{vmatrix}.$$

We denote the linear space of solutions of equation (1) as $S := \{u \in C^m[0, L] : \mathcal{L}u = 0\}$. Let us introduce new functions

$$\hat{v}^i[\mathbf{u}](x) := D(L_1, \dots, L_{i-1}, \delta_x, L_{i+1}, \dots, L_m)[\mathbf{u}], \quad i = \overline{1, m}. \quad (3)$$

The next lemma is valid.

Lemma 1 [see thesis, 1.4 lemma]. *Let $\{u^1, \dots, u^m\}$ be the basis of linear space S . Then the next propositions are equivalent:*


- a) *The functionals L_1, \dots, L_m are linearly independent;*
- b) *The functions $\hat{v}^1, \dots, \hat{v}^m$ are linearly independent;*
- c) *$D(\mathbf{L}) \neq 0$.*

If the functionals L_1, \dots, L_m are linearly independent, and


$$v^i(x) := \frac{D(L_1, \dots, L_{i-1}, \delta_x, L_{i+1}, \dots, L_m)[\mathbf{u}]}{D(\mathbf{L})[\mathbf{u}]} = \frac{\hat{v}^i[\mathbf{u}](x)}{D(\mathbf{L})[\mathbf{u}]}, \quad i = \overline{1, m},$$

then two bases $\{v^1, \dots, v^m\}$ and $\{L_1, \dots, L_m\}$ are biorthogonal: $\langle L_i, v^j \rangle = \delta_i^j$, $i, j = \overline{1, m}$.


Let us consider the finite set $\Xi_N = \{\xi_0, \xi_1, \dots, \xi_N\}$ where $\xi_0 = 0 < \xi_1 < \dots < \xi_{N-1} < \xi_N = L$ ($N \in \mathbb{N}$). The lines $y = \xi_i$, $i = \overline{0, N}$ as well as $y = x$ divide the square $[0, L]^2$ into triangles and trapezoids: D_x^i, D_y^i , $i = \overline{0, N-1}$ (see Fig. 1(b)). We use this



(a) classical fundamental solution



(b) generalized fundamental solution



(c)

Figure 1. Domains of the fundamental solutions

notation for triangles and trapezoids with boundary, i.e., D_x^i , D_y^i , $i = \overline{0, N-1}$ are closed sets.

Definition 1 [see thesis, 1.2 apibrėžimas]. The function $g(x, y)$ defined on $0 \leq x, y \leq L$ is called the *Generalized Fundamental Solution* (GFS) of homogeneous differential equation (1) if it has the following properties:

1. $g(x, y)$ has m partial derivatives in x in each figure D_y^i and D_x^i , $i = \overline{0, N-1}$ and these derivatives are continuous in both x and y ;
2. $g(x, y)$ (as a function of x) satisfies homogeneous equation (1) in each of those figures, i.e., $\mathcal{L}_x g(x, y) = 0$;
3. $g(x, y)$ is continuous in each rectangle $D_x^i \cup D_y^i$, $i = \overline{0, N-1}$ and has partial derivatives in x up to order $m-2$ and derivatives are continuous in both x and y in these rectangles;
4. The equality

$$\frac{\partial^{m-1} g(y+0, y)}{\partial x^{m-1}} - \frac{\partial^{m-1} g(y-0, y)}{\partial x^{m-1}} = 1$$

is valid for $y \in [0, L] \setminus \Xi_N$.

Definition 2 [see thesis, 1.3 apibrėžimas]. A Generalized Fundamental Solution $G(x, y)$ of equation (1) is called *Green's function* for problem (1)–(2) if it is satisfying (as a function of x) homogeneous additional conditions (2) for $y \in [0, L] \setminus \Xi_N$, i.e.,

$$\langle L_i(x), G(x, y) \rangle = 0, \quad i = 1, \dots, m, \quad y \in [0, L] \setminus \Xi_N.$$

If Green's function $G(x, y)$ exists for problem (1)–(2) then its solution allows the following integral representation:

$$u(x) = (G(x, y), f(y))_X = \int_0^L G(x, y) f(y) dy.$$

If functionals L_1, \dots, L_m are linearly independent then the following propositions about Green's functions are valid.

Lemma 2 [see thesis, 1.6 lemma]. *Green's function for differential equation (1) with homogeneous additional conditions $\langle L_1, u \rangle = 0, \dots, \langle L_m, u \rangle = 0$ is equal to:*

$$\begin{aligned} G(x, s) &= \langle \delta_x(y) - \mathbf{L}(y)\mathbf{v}(x), G^c(y, s) \rangle \\ &= G^c(x, s) - \sum_{i=1}^m \langle L_i(y), G^c(y, s) \rangle \frac{D(L_1, \dots, L_{i-1}, \delta_x, L_{i+1}, \dots, L_m)}{D(\mathbf{L})} \\ &= \frac{1}{D(\mathbf{L})[\mathbf{u}]} \begin{vmatrix} \langle L_1, u^1 \rangle & \dots & \langle L_m, u^1 \rangle & u^1(x) \\ \dots & \dots & \dots & \dots \\ \langle L_1, u^m \rangle & \dots & \langle L_m, u^m \rangle & u^m(x) \\ \langle L_1(\cdot), G^c(\cdot, s) \rangle & \dots & \langle L_m(\cdot), G^c(\cdot, s) \rangle & G^c(x, s) \end{vmatrix} \\ &= \frac{D(\mathbf{L}, \delta_x)[\mathbf{u}, G^c(\cdot, s)]}{D(\mathbf{L})[\mathbf{u}]} \end{aligned}$$

For the theoretical investigation of problems with nonlocal boundary conditions the next result about relations between Green's functions $G^u(x, s)$ and $G^v(x, s)$ of two inhomogeneous problems:

$$\begin{cases} \mathcal{L}u = f, \\ \langle l_i, u \rangle = 0, \quad i = 1, \dots, m, \end{cases} \quad \begin{cases} \mathcal{L}v = f, \\ \langle L_i, v \rangle = 0, \quad i = 1, \dots, m, \end{cases} \quad (4)$$

is useful.

Theorem 1 [see thesis, 1.2 teorema]. *The relations between Green's functions $G^v(x, s)$ and $G^u(x, s)$ for problems (4) are:*

$$\begin{aligned} G^v(x, s) &= \langle \delta_x(y) - \mathbf{L}(y)\mathbf{v}(x), G^u(y, s) \rangle \\ &= G^u(x, s) - \sum_{j=1}^m \langle L_j(y), G^u(y, s) \rangle \frac{D(L_1, \dots, L_{j-1}, \delta_x, L_{j+1}, \dots, L_m)}{D(\mathbf{L})} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{D(\mathbf{L})[\mathbf{u}]} \begin{vmatrix} \langle L_1, u^1 \rangle & \cdots & \langle L_m, u^1 \rangle & u^1(x) \\ \cdots & \cdots & \cdots & \cdots \\ \langle L_1, u^m \rangle & \cdots & \langle L_m, u^m \rangle & u^m(x) \\ \langle L_1(\cdot), G^u(\cdot, s) \rangle & \cdots & \langle L_m(\cdot), G^u(\cdot, s) \rangle & G^u(x, s) \end{vmatrix} \\
&= \frac{D(\mathbf{L}, \delta_x)[\mathbf{u}, G^u(\cdot, s)]}{D(\mathbf{L})[\mathbf{u}]}.
\end{aligned} \tag{5}$$

Let us investigate Green's function for the problem (1) with nonlocal boundary conditions

$$\mathcal{L}u := u^{(m)} + a^{m-1}(x)u^{(m-1)} + \cdots + a^1(x)u' + a^0(x)u = f(x), \tag{6}$$

$$\langle L_i, u \rangle := \langle \kappa_i, u \rangle - \gamma_i \langle \varkappa_i, u \rangle = 0, \quad i = \overline{1, m}. \tag{7}$$

We can write many problems with nonlocal boundary conditions (NBC) in this form, where $\langle \kappa_i, u \rangle := \langle \kappa_i(x), u(x) \rangle$ is a classical part, and $\langle \varkappa_i, u \rangle := \langle \varkappa_i(x), u(x) \rangle$, $i = \overline{1, m}$, is a nonlocal part of boundary conditions. For example, the functionals \varkappa_i , $i = \overline{1, m}$, can describe the multi-point ($\xi_j \in [0, 1]$, $j = \overline{1, m}$) or integral NBCs

$$\langle \varkappa, u \rangle = \sum_{j=1}^m \sum_{k=1}^m (\varkappa_j^k u^{(k-1)}(\xi_j)), \quad \langle \varkappa, u \rangle = \int_0^1 \varkappa(t)u(t) dt,$$

and the functionals κ_i , $i = \overline{1, m}$, can describe the local (classical) boundary conditions.

If $\gamma_1, \dots, \gamma_m = 0$, then problem (6)–(7) becomes classical. Suppose that there exists Green's function $G^{\text{cl}}(x, s)$ for this classical case. Then, Green's function exists for problem (6)–(7) if $\vartheta := D(\mathbf{L})[\mathbf{u}] \neq 0$. For $L_i = \kappa_i - \gamma_i \varkappa_i$, $i = \overline{1, m}$, we derive

$$\vartheta = \sum_{\sigma_1=0, \dots, \sigma_m=0}^1 \prod_{j=1}^m (-\gamma_j)^{\sigma_j} D((\varkappa_1^{\sigma_1} \kappa_1^{1-\sigma_1}), \dots, (\varkappa_m^{\sigma_m} \kappa_m^{1-\sigma_m})). \tag{8}$$

If we define the matrices $\mathbf{K} := (\kappa_{ij})$, $\kappa_{ij} = \langle \kappa_j, u_i \rangle$, $\mathbf{N} := (\varkappa_{ij})$, $\varkappa_{ij} = \langle \varkappa_j, u_i \rangle$, $\boldsymbol{\Gamma} := (\gamma_i \delta_{ij})$, then the condition $\vartheta \neq 0$ is equivalent to $\det(\mathbf{I} - \boldsymbol{\Gamma} \mathbf{N} \mathbf{K}^{-1}) \neq 0$. Since $\langle \kappa_i(\cdot), G^{\text{cl}}(\cdot, s) \rangle = 0$, $i = \overline{1, m}$, we can rewrite formula (5) as

$$G(x, s) = G^{\text{cl}}(x, s) + \sum_{j=1}^m \gamma_j \langle \varkappa_j(y), G^{\text{cl}}(y, s) \rangle v^j(x). \tag{9}$$

The main results of this section are published in [A3,A5,A8].

Chapter 2. The Application of Green's Function to the Second-Order Differential Problems with Nonlocal Boundary Conditions

In the second chapter the results of the first chapter are applied to the second-order differential problem:

$$-(p(x)u')' + q(x)u = f(x), \quad (10)$$

$$\langle L_1, u \rangle = g_1, \quad \langle L_2, u \rangle = g_2. \quad (11)$$

We investigate the differential equation $-u'' = f(x)$, $x \in (0, 1)$ with various type of nonlocal boundary conditions:

$$\begin{array}{lll} u(0) = \gamma_1 u(\xi_1), & u(1) = \gamma_2 u(\xi_2); & u'(0) = \gamma_1 u'(\xi_1), & u(1) = \gamma_2 u(\xi_2); \\ u(0) = \gamma_1 u(\xi_1), & u(1) = \gamma_2 u'(\xi_2); & u'(0) = \gamma_1 u(\xi_1), & u(1) = \gamma_2 u'(\xi_2); \\ u(0) = \gamma_1 u'(\xi_1), & u(1) = \gamma_2 u(\xi_2); & u'(0) = \gamma_1 u'(\xi_1), & u(1) = \gamma_2 u(\xi_2); \\ u(0) = \gamma_1 u'(\xi_1), & u(1) = \gamma_2 u'(\xi_2); & u'(0) = \gamma_1 u'(\xi_1), & u(1) = \gamma_2 u'(\xi_2); \\ u(0) = \gamma_1 u(\xi_1), & u'(1) = \gamma_2 u(\xi_2); & u'(0) = \gamma_1 u(\xi_1), & u'(1) = \gamma_2 u'(\xi_2); \\ u(0) = \gamma_1 u(\xi_1), & u'(1) = \gamma_2 u'(\xi_2); & u'(0) = \gamma_1 u'(\xi_1), & u'(1) = \gamma_2 u(\xi_2); \\ u(0) = \gamma_1 u'(\xi_1), & u'(1) = \gamma_2 u(\xi_2); & u'(0) = \gamma_1 u(\xi_1), & u'(1) = \gamma_2 u(\xi_2); \\ u(0) = \gamma_1 u'(\xi_1), & u'(1) = \gamma_2 u'(\xi_2); & u'(0) = \gamma_1 u'(\xi_1), & u'(1) = \gamma_2 u'(\xi_2); \end{array}$$

or with integral nonlocal boundary conditions

$$u(0) = \gamma_1 \int_0^1 \alpha_1(x)u(x) dx, \quad u(1) = \gamma_2 \int_0^1 \alpha_2(x)u(x) dx,$$

where $\alpha_1, \alpha_2 \in L^1(0, 1)$. Also, the examples with the differential operators $\mathcal{L}u := -u'' + u$, $\mathcal{L}u := -u'' - u$, $\mathcal{L}u := -u'' - \frac{2x}{1+x^2}u'$ are presented.

For these problems we can find the fundamental system for homogeneous equation. In all cases we find Green's function, expressing it either via Green's function for problem with initial conditions (see Lemma 2), or via Green's function for problem with classical boundary conditions, when $\gamma_1 = \gamma_2 = 0$ (see formula (9)).

The main results of this section are published in [A1,A2,B1].

Chapter 3. The Second-Order Discrete Problem

The differential equation of the second chapter can be approximated by the difference equation

$$\mathcal{L}u := a_i^2 u_{i+2} + a_i^1 u_{i+1} + a_i^0 u_i = f_i, \quad (12)$$

where $a^2, a^0 \neq 0$, which is investigated in the third chapter.

Let $X = \{0, 1, \dots, n\}$, $\tilde{X} = \{0, 1, \dots, n-2\}$. In discrete case, the notation and definition are as follows:

$$\begin{aligned} [\mathbf{u}]_{ij} &= [u^1, u^2]_{ij} := \begin{vmatrix} u_i^1 & u_j^1 \\ u_i^2 & u_j^2 \end{vmatrix}, \quad D[\mathbf{u}]_{ij} = \det[u^1, u^2]_{ij} := \begin{vmatrix} u_i^1 & u_j^1 \\ u_i^2 & u_j^2 \end{vmatrix}, \\ W[\mathbf{u}]_j &:= \begin{vmatrix} u_{j-1}^1 & u_{j-1}^2 \\ u_j^1 & u_j^2 \end{vmatrix} = \begin{vmatrix} u_{j-1}^1 & u_j^1 \\ u_{j-1}^2 & u_j^2 \end{vmatrix} = D[\mathbf{u}]_{j-1,j}, \quad j = 1, \dots, n, \\ V[\mathbf{u}]_{ij} &:= \frac{D[\mathbf{u}]_{j+1,i}}{W[\mathbf{u}]_{j+2}} = \frac{D[\mathbf{u}]_{j+1,i}}{D[\mathbf{u}]_{j+1,j+2}}, \quad i \in X, \quad j = -1, 0, 1, \dots, n-2, \end{aligned}$$

when $W[\mathbf{u}]_{j+2} \neq 0$. We define $V[\mathbf{u}]_{i,n-1} = V[\mathbf{u}]_{in} = 0$, $i \in X$. Note that $V[\mathbf{u}]_{j+1,j} = 0$, $V[\mathbf{u}]_{j+2,j} = 1$ for $j \in \tilde{X}$. For the functionals $\mathbf{f} = (f_1, f_2)$ and the functions $\mathbf{w} = [w^1, w^2]$, we define the determinant:

$$D(\mathbf{f})[\mathbf{w}] := \begin{vmatrix} \langle f_1, w^1 \rangle & \langle f_2, w^1 \rangle \\ \langle f_1, w^2 \rangle & \langle f_2, w^2 \rangle \end{vmatrix}.$$

We introduce a function $G^c \in F(X \times \tilde{X})$:

$$G_{ij}^c := H_{i-j} V_{ij} / a_j^2, \quad \text{where } H_i := \begin{cases} 1, & i > 0, \\ 0, & i \leq 0. \end{cases}$$

We consider the solutions of homogeneous equation (12). Let $S \subset F(X)$ be a two-dimensional linear space of solutions. We investigate additional equations:

$$\langle L_1, u \rangle = 0, \quad \langle L_2, u \rangle = 0, \quad u \in S, \quad (13)$$

where L_1, L_2 are linearly independent linear functionals and we use the notation $\mathbf{L} = (L_1, L_2)$. We introduce new functions

$$\bar{v}_i^1 := D(\delta_i, L_2)[\mathbf{u}], \quad \bar{v}_i^2 := D(L_1, \delta_i)[\mathbf{u}]. \quad (14)$$

Lemma 3 [see thesis, 3.3 lemma]. Let $\{u^1, u^2\}$ be the basis of the linear space S . Then the following propositions are equivalent:

- a) The functionals L_1, L_2 are linearly independent;
- b) The functions \bar{v}^1, \bar{v}^2 are linearly independent;
- c) $D(\mathbf{L})[\mathbf{u}] \neq 0$.

Suppose that $X_n := X = \{0, 1, \dots, n\}$, $F(X) := \{u \mid u : X \rightarrow \mathbb{K}\}$ be a linear space of real (complex) functions. Let $A : F(X_n) \rightarrow F(X_{n-m}) = \text{Im } A$ and $B : F(X_n) \rightarrow F(X_{M-n+m})$ be linear operators, $0 \leq m \leq n$. Consider the operator equation $Au = f$, where $u \in F(X_n)$ is unknown and $f \in F(X_{n-m})$ is given, with the additional operator equation $Bu = 0$.

If the solution of the problem $Au = f, Bu = 0$ allows the following representation:

$$u_i = \sum_{j=0}^{n-m} G_{ij} f_j, \quad i \in X_n,$$

then $G \in F(X_n \times X_{n-m})$ is called *discrete Green's function* of operator A with the additional condition $Bu = 0$. Green's function exists, if $\text{Ker } A \cap \text{Ker } B = \{0\}$. For $m = 2$, $u_i = \sum_{j=0}^{n-2} G_{ij} f_j$, $i \in X_n$.

Lemma 4 [see thesis, 3.5 lemma]. *Green's function for problem (12) with homogeneous additional conditions $\langle L_1, u \rangle = 0, \langle L_2, u \rangle = 0$, where the functionals L_1 and L_2 are linearly independent, is equal to:*

$$G_{ij} = \frac{D(\mathbf{L}, \delta_i)[\mathbf{u}, G_{\cdot,j}^c]}{D(\mathbf{L})[\mathbf{u}]}, \quad i \in X, j \in \tilde{X}.$$

For the theoretical investigation of problems with NBCs, the next result about the relations between Green's functions G_{ij}^u and G_{ij}^v of two nonhomogeneous problems

$$\begin{cases} \mathcal{L}u = f, \\ \langle l_m, u \rangle = 0, \quad m = 1, 2, \end{cases} \quad \begin{cases} \mathcal{L}v = f, \\ \langle l_m, v \rangle = 0, \quad m = 1, 2, \end{cases} \quad (15)$$

is useful.

Theorem 2 [see thesis, 3.2 teorema]. *If Green's function G^u exists and functionals L_1 and L_2 are linearly independent, then*

$$G_{ij}^v = \frac{D(\mathbf{L}, \delta_i)[\mathbf{u}, G_{\cdot,j}^u]}{D(\mathbf{L})[\mathbf{u}]}, \quad i \in X, j \in \tilde{X}. \quad (16)$$

Let us investigate Green's function for the problem with nonlocal boundary conditions

$$\mathcal{L}u := a_i^2 u_{i+2} + a_i^1 u_{i+1} + a_i^0 u_i = f_i, \quad i \in \tilde{X}, \quad (17)$$

$$\langle L_1, u \rangle := \langle \kappa_1, u \rangle - \gamma_1 \langle \varkappa_1, u \rangle = 0, \quad (18)$$

$$\langle L_2, u \rangle := \langle \kappa_2, u \rangle - \gamma_2 \langle \varkappa_2, u \rangle = 0. \quad (19)$$

We can write many problems with nonlocal boundary conditions (NBC) in this form, where $\langle \kappa_m, u \rangle := \langle \kappa_m^i, u_i \rangle$, $m = 1, 2$, is a classical part, and $\langle \varkappa_m, u \rangle := \langle \varkappa_m^i, u_i \rangle$, $m = 1, 2$, is a nonlocal part of boundary conditions.

If $\gamma_1, \gamma_2 = 0$, then problem (17)–(19) becomes classical. Suppose that there exists Green's function G_{ij}^{cl} for the classical case. Then Green's function exists for problem (17)–(19), if $\vartheta := D(\mathbf{L})[\mathbf{u}] \neq 0$. For $L_m = \kappa_m - \gamma_m \varkappa_m$, $m = 1, 2$, we derive

$$\vartheta = D(\kappa_1 \cdot \kappa_2)[\mathbf{u}] - \gamma_1 D(\varkappa_1 \cdot \kappa_2)[\mathbf{u}] - \gamma_2 D(\kappa_1 \cdot \varkappa_2)[\mathbf{u}] + \gamma_1 \gamma_2 D(\varkappa_1 \cdot \varkappa_2)[\mathbf{u}].$$

Since $\langle \kappa_m^k, G_{kj}^{\text{cl}} \rangle = 0$, $m = 1, 2$, we can rewrite formula (16) as

$$\begin{aligned} G_{ij} &= G_{ij}^{\text{cl}} + \gamma_1 \langle \varkappa_1^k, G_{kj}^{\text{cl}} \rangle \frac{D(\delta_i, L_2)}{\vartheta} + \gamma_2 \langle \varkappa_2^k, G_{kj}^{\text{cl}} \rangle \frac{D(L_1, \delta_i)}{\vartheta} \\ &= \frac{1}{\vartheta} \begin{vmatrix} \langle L_1, u^1 \rangle & \langle L_2, u^1 \rangle & u_i^1 \\ \langle L_1, u^2 \rangle & \langle L_2, u^2 \rangle & u_i^2 \\ -\gamma_1 \langle \varkappa_1^k, G_{kj}^{\text{cl}} \rangle & -\gamma_2 \langle \varkappa_2^k, G_{kj}^{\text{cl}} \rangle & G_{ij}^{\text{cl}} \end{vmatrix}. \end{aligned} \quad (20)$$

The main results of this section are published in [A4].

Chapter 4. The m -Order Discrete Problem

In this chapter the results of the third chapter are generalized. We consider the m -order discrete problem with m additional conditions

$$a_i^m u_{i+m} + \cdots + a_i^2 u_{i+2} + a_i^1 u_{i+1} + a_i^0 u_i = f_i, \quad i \in \tilde{X} = \{0, 1, \dots, n-m\}, \quad (21)$$

$$\langle L_1, u \rangle = g_1 \in \mathbb{K}, \quad \dots, \quad \langle L_m, u \rangle = g_m \in \mathbb{K}. \quad (22)$$

The expression of the solution and Green's function for this problem are formulated; also the relation between two Green's functions for two problems with the same equation, but with various linear conditions, is given.

Let $\mathbf{L} = (L_1, \dots, L_m)$. We consider a determinant $D[\mathbf{u}]_i : X^m \rightarrow \mathbb{K}$:

$$D[\mathbf{u}]_i = \det[\mathbf{u}]_i = \det[u^1, \dots, u^m]_{i_1 \dots i_m} := \begin{vmatrix} u_{i_1}^1 & \cdots & u_{i_m}^1 \\ \cdots & \cdots & \cdots \\ u_{i_1}^m & \cdots & u_{i_m}^m \end{vmatrix}.$$

The Wronskian determinant $W[\mathbf{u}]_j$ and similar determinant $\tilde{W}[\mathbf{u}]_{ij}$ in the theory of difference equations are denoted as follows

$$W[\mathbf{u}]_j := \begin{vmatrix} u_{j-m+1}^1 & \cdots & u_{j-m+1}^m \\ \cdots & \cdots & \cdots \\ u_{j-1}^1 & \cdots & u_{j-1}^m \\ u_j^1 & \cdots & u_j^m \end{vmatrix} = D[\mathbf{u}]_{j-m+1, \dots, j}, \quad j = m-1, \dots, n,$$

$$\tilde{W}[\mathbf{u}]_{ij} := \begin{vmatrix} u_{j-m+1}^1 & \cdots & u_{j-m+1}^m \\ \cdots & \cdots & \cdots \\ u_{j-1}^1 & \cdots & u_{j-1}^m \\ u_i^1 & \cdots & u_i^m \end{vmatrix}, \quad i \in X, \quad j = m-1, \dots, n+1.$$

Let (if $W[\mathbf{u}]_{j+m} \neq 0$)

$$V[\mathbf{u}]_{ij} := \frac{\tilde{W}[\mathbf{u}]_{i,j+m}}{W[\mathbf{u}]_{j+m}}, \quad i \in X, \quad j \in -1, \dots, n-m.$$

We define $V[\mathbf{u}]_{i,n-k} = 0$, $k = 0, \dots, m-1$, $i \in X$. Note that $V[\mathbf{u}]_{j+k,j} = 0$, $k = 1, \dots, m-1$, $V[\mathbf{u}]_{j+m,j} = 1$ for $j \in \tilde{X}$.

We introduce a function $G^c \in F(X \times \tilde{X})$:

$$G_{ij}^c := H_{i-j} V_{ij} / a_j^m.$$

Lemma 5 [see thesis, 4.2 lemma]. *Green's function for problem (21) with the homogeneous additional conditions $\langle L_1, u \rangle = 0, \dots, \langle L_m, u \rangle = 0$, where functionals L_1, \dots, L_m are linearly independent, is equal to*

$$G_{ij} = \frac{D(\mathbf{L}, \delta_i)[\mathbf{u}, G_{\cdot,j}^c]}{D(\mathbf{L})[\mathbf{u}]}, \quad i \in X, \quad j \in \tilde{X}. \quad (23)$$

For the theoretical investigation of problems with NBCs, the next result about the relations between Green's functions G_{ij}^u and G_{ij}^v of two nonhomogeneous problems

$$\begin{cases} \mathcal{L}u = f, \\ \langle l_k, u \rangle = 0, \quad k = 1, \dots, m, \end{cases} \quad \begin{cases} \mathcal{L}v = f, \\ \langle L_k, v \rangle = 0, \quad k = 1, \dots, m, \end{cases} \quad (24)$$

is useful.

Theorem 3 [see thesis, 4.2 teorema]. *If Green's function G^u exists and the functionals L_1, \dots, L_m are linearly independent, then*

$$\begin{aligned} G_{ij}^v &= \frac{D(\mathbf{L}, \delta_i)[\mathbf{u}, G_{:,j}^u]}{D(\mathbf{L})[\mathbf{u}]} \\ &= \frac{1}{D(\mathbf{L})[\mathbf{u}]} \begin{vmatrix} \langle L_1, u^1 \rangle & \cdots & \langle L_m, u^1 \rangle & u_i^1 \\ \cdots & \cdots & \cdots & \cdots \\ \langle L_1, u^m \rangle & \cdots & \langle L_m, u^m \rangle & u_i^m \\ \langle L_1^k, G_{kj}^u \rangle & \cdots & \langle L_m^k, G_{kj}^u \rangle & G_{ij}^u \end{vmatrix} \\ &= G_{ij}^u - \langle L_1^k, G_{kj}^u \rangle \frac{D(\delta_i, L_2, \dots, L_m)}{D(\mathbf{L})} - \dots \\ &\quad - \langle L_m^k, G_{kj}^u \rangle \frac{D(L_1, \dots, L_{m-1}, \delta_i)}{D(\mathbf{L})}, \quad i \in X, j \in \tilde{X}. \end{aligned} \quad (25)$$

Let us investigate Green's function for the problem with nonlocal boundary conditions

$$\mathcal{L}u := a_i^m u_{i+m} + \dots + a_i^1 u_{i+1} + a_i^0 u_i = f_i, \quad i \in \tilde{X}, \quad (26)$$

$$\langle L_l, u \rangle := \langle \kappa_l, u \rangle - \gamma_l \langle \varkappa_l, u \rangle = 0, \quad l = \overline{1, m}. \quad (27)$$

We can write many problems with nonlocal boundary conditions (NBC) in this form, where $\langle \kappa_l, u \rangle := \langle \kappa_l^i, u_i \rangle$, $l = \overline{1, m}$, is a classical part, and $\langle \varkappa_l, u \rangle := \langle \varkappa_l^i, u_i \rangle$, $l = \overline{1, m}$, is a nonlocal part of boundary conditions (BC).

If $\gamma_1, \dots, \gamma_m = 0$, then problem (26)–(27) becomes classical. Suppose that there exists Green's function G_{ij}^{cl} for the classical case. Then Green's function exists for problem (26)–(27), if $\vartheta := D(\mathbf{L})[\mathbf{u}] \neq 0$. For $L_l = \kappa_l - \gamma_l \varkappa_l$, $l = \overline{1, m}$, we derive that

$$\vartheta = \sum_{j_1, \dots, j_m=0}^1 \gamma_1^{j_1} \cdots \gamma_m^{j_m} D((1-j_1)\kappa_1 - j_1 \varkappa_1, \dots, ((1-j_m)\kappa_m - j_m \varkappa_m))[\mathbf{u}].$$

Since $\langle \kappa_l^k, G_{kj}^{\text{cl}} \rangle = 0$, $l = \overline{1, m}$, we can rewrite formula (25) as follows:

$$\begin{aligned}
G_{ij} &= G_{ij}^{\text{cl}} + \gamma_1 \langle \varkappa_1^k, G_{kj}^{\text{cl}} \rangle \frac{D(\delta_i, L_2, \dots, L_m)}{\vartheta} + \dots \\
&\quad + \gamma_m \langle \varkappa_m^k, G_{kj}^{\text{cl}} \rangle \frac{D(L_1, \dots, L_{m-1}, \delta_i)}{\vartheta} \\
&= \frac{1}{\vartheta} \begin{vmatrix} \langle L_1, u^1 \rangle & \dots & \langle L_m, u^1 \rangle & u_i^1 \\ \dots & \dots & \dots & \dots \\ \langle L_1, u^m \rangle & \dots & \langle L_m, u^m \rangle & u_i^m \\ -\gamma_1 \langle \varkappa_1^k, G_{kj}^{\text{cl}} \rangle & \dots & -\gamma_m \langle \varkappa_m^k, G_{kj}^{\text{cl}} \rangle & G_{ij}^{\text{cl}} \end{vmatrix}. \tag{28}
\end{aligned}$$

The main results of this section are presented in [A9].

General conclusions

1. The m -order linear nonhomogeneous differential and discrete equations with various conditions are analyzed. The expressions of the solutions and Green's functions for these problems are obtained.
2. The independence of linear functionals is necessary and sufficient condition for the existence of Green's function for the problem which conditions are written with linear functionals.
3. Green's functions for problems with the same equation are related to each other. If the fundamental system of homogeneous equation and Green's function are known, then Green's function of the other problem with the same equation can be expressed via Green's function of the first problem.
4. The formulae can be applied to a very wide class of problems with nonconstant coefficients and various boundary conditions as well as nonlocal boundary conditions.
5. The expressions and relations for Green's function are the same in both differential and difference cases. The conditions of the existence of Green's function are also formulated in the same way.

List of Published Works on the Topic of the Dissertation

In the reviewed scientific periodical publications

- [A1] S. Roman and A. Štikonas. Green's functions for stationary problems with nonlocal boundary conditions. *Lith. Math. J.*, **49**(2):190–202, 2009.
Doi:10.1007/s10986-009-9041-0.
- [A2] A. Štikonas and S. Roman. Stationary problems with two additional conditions and formulae for Green's functions. *Numer. Funct. Anal. Optim.*, **30**(9):1125–1144, 2009.
Doi:10.1080/01630560903420932.
- [A3] S. Roman and A. Štikonas. Third-order linear differential equation with three additional conditions and formula for Green's function. *Lith. Math. J.*, **50**(4):426–446, 2010.
Doi:10.1007/s10986-010-9097-x.
- [A4] S. Roman and A. Štikonas. Green's function for discrete second-order problems with nonlocal boundary conditions. *Bound. Value Probl.*, **2011**:1–23, 2011.
Doi:10.1155/2011/767024.
- [A5] S. Roman. Linear differential equation with additional conditions and formulae for Green's function. *Math. Model. Anal.*, **16**(3):401–417, 2011.
Doi:10.3846/13926292.2011.602125.
- [A6] S. Roman and A. Štikonas. Nelokaliųjų stacionariųjų kraštinių uždavinių Gryno funkcijos. *Liet. mat. rink. LMD darbai*, **48/49**:333–337, 2008.
- [A7] S. Roman and A. Štikonas. The properties of Green's functions for one stationary problem with nonlocal boundary conditions. *Liet. mat. rink. LMD darbai*, **50**:340–344, 2009.
- [A8] S. Roman and A. Štikonas. Linear ODE with nonlocal boundary conditions and Green's functions for such problems. *Liet. mat. rink. LMD darbai*, **51**:379–384, 2010.
- [A9] S. Roman and A. Štikonas. Green's function for discrete problems with nonlocal boundary conditions. *Liet. mat. rink. LMD darbai*, **52**, 2011. (in press)

In the other editions

- [B1] S. Roman and A. Štikonas. Green's functions for stationary problems with four-point non-local boundary conditions. In V. Kleiza, S. Rutkauskas and A. Štikonas(Eds.), *Differential Equations and Their Applications (DETA '2009)*, pp. 123–130. Kaunas University of Technology, 2009.

About the author

Svetlana Roman was born in Trakai on 21 of July 1983.

First degree in Mathematics, Faculty of Mathematics and Informatics, Vilnius University, 2005. Master of Science in Mathematics, Faculty of Mathematics and Informatics, Vilnius University, 2007. In 2007–2011 – PhD student of Institute of Mathematics and Informatics. 2007 – at present – Copyeditor of scientific articles in Ltd. “VTEX”.

GRYNO FUNKCIJOS UŽDAVINIAMS SU NELOKALIOSIOMIS KRAŠTINĖMIS SĀLYGOMIS

Problemos formulavimas

Disertacijoje tiriamos antros ir m -tosios eilės diferencialinių ir diskrečiųjų lygčių su įvairiomis sąlygomis, kurios yra aprašytos tiesiškai nepriklausomais tiesiniais funkcionalais, *Gryno funkcijos*. Didelis dėmesys skirtas uždaviniams su nelokaliosiomis kraštinėmis sąlygomis.

Darbo aktualumas

Diferencialinių lygčių teorijoje pagrindinės sąvokos buvo suformuluotos tiriant klasikines matematinės fizikos uždavinius. Tačiau šiuolaikinės problemos skatina naujų uždavinijų formulavimą ir tyrimą, pavyzdžiui, nelokaliųjų uždavinių klasę. Nelokaliosios sąlygos atsiranda tada, kai negalima tiesiogiai išmatuoti duomenų nagrinėjamo uždavinio srities krašte. Šiuo atveju formuluojami uždaviniai, kuriuose sprendinio ir / arba jo išvestinės reikšmės susijusios keliuose taškuose ar ištisame intervale.

1963 m. J. R. Kenonas nagrinėjo uždavinį, kuris dabar vadinamas *nelokaliuoju* (terminą *nelokaliosios kraštinės sąlygos*, pirmą sykį panaudojo N. I. Jonkinas 1977 m.). Panašū uždavinį tyrė L. I. Kamyninas 1964 m. Vieni iš pirmųjų nelokaliuosius krašinius uždavinius 1969 m. nagrinėjo A. A. Samarskis ir A. V. Bitsadzė. Jų vardais pavadintos tam tikros nelokaliosios sąlygos vienamačiu atveju (t. y. Bitsadzės ir Samarskio sąlygos). Diferencialines lygtis (pavyzdžiui, paprastąją, elipsinę, parabolinę, ir kitas) su įvairaus tipo nelokaliosiomis sąlygomis nagrinėjo mokslinginkai: Iljinė, Moisejevas, Jonkinas, Gulinas, Vebas, Infantė ir kt. Lietuvoje tokius uždavinius tyrė Sapagovas, Čiegiš, Štikonas ir jų mokiniai.

Šioje disertacijoje tiriami antros ir aukštesnės eilės diferencialinių ir diskrečiųjų lygtių su įvairiomis sąlygomis (pavyzdžiu, su pradinėmis, kraštinėmis arba nelokaliosiomis sąlygomis) Gryno funkcijos.

Matematikoje Gryno funkcija naudojama sprendžiant nehomogenines lygtis atsižvelgiant į kraštines sąlygas. Tiesinių paprastųjų ir dalinių išvestinių diferencialinių lygtių teorijoje Gryno funkcija atlieka svarbų vaidmenį. Jos pagalba galima rasti stacionariųjų ir nestacionariųjų uždavinių sprendinius, ištirti sprendinių egzistavimą ir vienatį. Gryno funkcijos taikomos elektrostatikoje (Puasono lygties sprendimas), kondensuotųjų medžiagų teorijoje (difuzijos lygties sprendimas), kvantinėje mechanikoje (Šredingerio lygties sprendimas).

Uždavinių su klasikinėmis kraštinėmis sąlygomis Gryno funkcijas nagrinėjo V. S. Vladimirovas, U. M. Ašeris, R. M. M. Matteijus, R. D. Raselas, I. Stakgoldas, D. G. Dafis. Be to, A. A. Samarskis, E. S. Nikolajevas ir A. V. Gulinas tyrė diskrečiųjų uždavinių Gryno funkcijas.

Pusiau tiesinių uždavinių (dešinioji diferencialinės lygties pusė netiesiškai priklauso nuo ieškomosios funkcijos, o kairioji yra tiesinis diferencialinis operatorius) su nelokaliosiomis kraštinėmis sąlygomis tyrimas ir teigiamų sprendinių egzistavimas remiasi tiesinių uždavinių su nelokaliosiomis kraštinėmis sąlygomis Gryno funkcijos tyrimu.

Antros ir aukštesnės eilės kraštinių uždavinių su įvairiomis nelokaliosiomis kraštinėmis sąlygomis Gryno funkcijas konstravo D. R. Andersonas, G. Infanté, L. X. Truongas, J. R. L. Vebas, B. Jangas, Z. Zao ir kiti mokslininkai. Šiuose darbuose autoriai nagrinėjo sprendinių egzistavimą ir vienatį, naudodami įvairus metodus: apatinio ir viršutinio sprendimo metodą; Krasnoselskio nejudamojo taško teoremą; Averio ir Petersono nejudamojo taško teoremą; Leréjaus ir Šauderio pratešimo principą; Legeto ir Viljamso nejudamojo taško teoremą.

Šios disertacijos *pirmajame* skyriuje nagrinėjama m -tosios eilės tiesinė diferencialinė lygtis su įvairiomis sąlygomis, kurios gali būti ne tik klasikinės, bet ir nelokaliosios. Šios sąlygos formuluojamos nepriklausomų tiesinių funkcialų pagalba. Taip pat pirmajame skyriuje išvesta formulė, apibendrinanti uždavinių su neklasikinėmis ir klasikinėmis kraštinėmis sąlygomis sąryšį.

Antrajame skyriuje suformuluota antros eilės diferencialinė lygtis. Pirmojo skyriaus rezultatai pritaikomi šiam uždaviniui. Pateikiami pavyzdžiai, kuriuose pavaizduota, kaip

konstruojama uždavinių su įvairiomis nelokaliosiomis kraštinėmis sąlygomis Gryno funkcija.

Pirmojo ir antrojo skyriaus nagrinėjamas diferencialines lygtis galima aproksimuoti baigtinių skirtumų lygtimis. *Trečiąjame* skyriuje tiriama antros eilės diskrečioji lygtis su dviem sąlygomis.

Ketvirtąjame disertacijos skyriuje apibendrinti trečiojo skyriaus rezultatai m -tosios eilės diskrečiajai lygčiai su įvairiomis sąlygomis.

Tyrimų objektas

Disertacijos tyrimo objektas – diferencialinės ir diskrečiosios lygtys su įvairiomis sąlygomis, šiu uždavinių sprendiniai ir Gryno funkcijos, jų egzistavimo sąlygos ir rezultatų pritaikymas uždaviniams su nelokaliosiomis kraštinėmis sąlygomis.

Darbo tikslas ir uždaviniai

Disertacijos tikslas – ištirti tiesinius diferencialinių ir diskretūjų uždavinius su įvairiomis sąlygomis, užrašyti nehomogeninės lygties sprendinių išraiškas, kai žinoma fundamentalioji sprendinių sistema, rasti Gryno funkcijas, taip pat nustatyti saryši tarp dviejų Gryno funkcijų uždaviniams su ta pačia lygtimi ir skirtingomis sąlygomis.

Siekiant numatyto tikslo buvo sprendžiami šie uždaviniai:

- ištirti homogeninės lygties sprendinių tiesinę erdvę;
- ištirti nehomogeninių diferencialinio ir diskrečiojo uždavinių sprendinius su ta pačia lygtimi, bet su skirtingomis sąlygomis;
- surasti diferencialinio ir diskrečiojo uždavinių Gryno funkcijas ir jų egzistavimo sąlygas;
- ištirti dviejų diferencialinių arba diskrečiųjų lygčių su skirtingomis sąlygomis Gryno funkcijas;
- pritaikyti gautus rezultatus uždaviniams su nelokaliosiomis kraštinėmis sąlygomis;
- palyginti diferencialinio ir diskrečiojo uždavinių rezultatus.

Tyrimų metodika

Disertacijos pagrindinių formulų išvedimui buvo taikomas konstantų varijavimo metodas. Naudojant šį metodą ir determinantų savybes, galima surasti Gryno funkcijas diferencialinėms ir diskrečiosioms lygtims su kintamais koeficientais ir įvairiomis kraštinėmis sąlygomis. Taip pat darbe aprašyti tiesiniai funkcionalai ir funkcionaliniai determinantai, kurių savybėmis buvo naudojamas. Braižant grafikus, buvo naudojamas „MAPLE“ programų paketas.

Darbo mokslinis naujumas ir jo reikšmė

Tiriant antros ir aukštesnės eilės (tieki tiesinių, tieki netiesinių) uždavinių sprendinių egzistavimą ir vienatį, dažniausiai konstruojama Gryno funkcija tiesiniam uždavinui. Beveik visais atvejais Gryno funkcija ieškoma uždavinui, kuris yra šios disertacijos uždavinio atskiras atvejis.

Šioje disertacijoje nagrinėjamos Gryno funkcijos uždaviniams su bet kokiomis tiesinėmis sąlygomis. Rasta būtina ir pakankama Gryno funkcijos egzistavimo sąlyga. Taip pat gautas Gryno funkcijų sąryšis, kuris padeda ištirti Gryno funkciją uždavinui su nelokaliosiomis kraštinėmis sąlygomis, žinant Gryno funkciją uždavinui su pradinėmis arba klasikinėmis kraštinėmis sąlygomis. Dažniausiai įvairose monografijose ir straipsniuose tokio tipo formulės buvo gautos uždavinui su klasikinėmis kraštinėmis sąlygomis, bet šiame darbe jos yra išvestos uždavinui su bet kokiomis (tiesinėmis) sąlygomis. Tokiu būdu, mes galime užrašyti Gryno funkcijos išraiška uždavinui su įvairiomis (pavyzdžiui, nelokaliosiomis) sąlygomis, jei žinomas homogeninės lygties fundamentalioji bazė ir uždavinio su kitomis (pavyzdžiui, klasikinėmis) sąlygomis Gryno funkcija. Gautus rezultatus galima pritaikyti uždaviniams su nelokaliosiomis kraštinėmis sąlygomis (ir ne tik tokiomis).

Darbo rezultatų praktinė reikšmė

Disertacijos gauti rezultatai gali būti panaudojami sprendžiant tiesines diferencialines arba diskrečiasias lygtis su įvairiomis (pavyzdžiui, pradinėmis, kraštinėmis, nelokaliosiomis) tiesinėmis sąlygomis, o taip pat tiriant sprendinių egzistavimą, vienatį ir jų savybes.

Ginamieji teiginiai

- Būtina ir pakankama Gryno funkcijos egzistavimo sąlyga.
- Antros ir m -tosios eilės diferencialinio ir diskrečiojo uždavinių su įvairiomis tiesinėmis sąlygomis Gryno funkcijų išraiškos.
- Gryno funkcijos sąryšis su Gryno funkcija uždaviniui su pradinėmis sąlygomis.
- Uždavinių su ta pačia lygtimi, bet su skirtingomis sąlygomis, Gryno funkcijų sąryšis.
- Taikymai uždaviniams su nelokaliomis kraštinėmis sąlygomis.

Disertacijos struktūra

Disertaciją sudaro įvadas, 4 skyriai, išvados, literatūros sąrašas ir autorės publikacijų disertacijos tema sąrašas. Bendra disertacijos apimtis – 129 puslapiai, 4 grafikai, 4 lentelės. Disertacijos rezultatai skelbti 9 publikacijose. 1 publikacija priimta spaudai Lietuvos periodiniame recenzuojamame leidinyje. Šia tema skaityta 9 pranėsimai mokslinėse konferencijose.

Šios disertacijos *pirmajame skyriuje* nagrinėjama m -tosios eilės diferencialinė lygtis su įvairiomis (iskaitant ir nelokaliomis) sąlygomis. *Antrajame skyriuje* pirmojo skyriaus rezultatai pritaikyti antros eilės diferencialiniams uždaviniui. Pateikiami pavyzdžiai, kuriuose demonstruojama kaip konstruoti Gryno funkciją uždaviniams su įvairiomis nelokaliomis kraštinėmis sąlygomis. Antros eilės diferencialines lygtis galima aproksimuoti diskrečiosiomis lygtimis, kurios nagrinėjamos *trečiąjame skyriuje*. Trečiojo skyriaus rezultatai apibendrinti *ketvirtajame skyriuje*. Šiame skyriuje nagrinėjamas m -tosios eilės diskretusis uždavinys su įvairiomis sąlygomis.

Bendrosios išvados

1. Išanalizuotos m -tosios eilės tiesinės nehomogeninės diferencialinė ir diskrečioji lygtys su įvairiomis sąlygomis. Gautos šių uždavinių sprendinių išraiškos ir Gryno funkcijos.
2. Tiesinio nehomogeninio uždavinio, kurio sąlygos užrašomas tiesiniais funkcionais, būtina ir pakankama Gryno funkcijos egzistavimo sąlyga yra tų funkcionalų tiesinis nepriklausomumas.

3. Gryno funkcijos uždaviniam su ta pačia lygtimi yra susietos sąryšiu. Jei žinome vieną Gryno funkciją ir homogeninės lyties fundamentaliąją sistemą, tai kito uždavinio su ta pačia lygtimi, Gryno funkciją galima išreikšti per pirmojo uždavinio Gryno funkciją.

4. Formulės gali būti pritaikytos plačiai uždaviniių klasei su nepastoviais koeficientais ir įvairiomis kraštinėmis sąlygomis, tame tarpe su nelokaliosiomis kraštinėmis sąlygomis.

5. Gautos Gryno funkcijų formulės yra vienodos tiek diferencialiniu, tiek diskrečiuoju atvejais. Gryno funkcijų egzistavimo sąlygos irgi formuluojamos vienodai.

Trumpos žinios apie autore

Svetlana Roman gimė 1983 m. liepos 21 d. Trakuose. 2005 m. įgijo matematikos bacheloro laipsnį Vilniaus universitete Matematikos ir informatikos fakultete. 2007 m. baigė Matematinio modeliavimo ir optimizavimo magistrantūros studijų programą ir įgijo matematikos magistro kvalifikaciją laipsnį Vilniaus universitete Matematikos ir informatikos fakultete. 2007–2011 m. studijavo matematikos krypties doktorantūroje Vilniaus universiteto Matematikos ir informatikos institute. Nuo 2007 m. dirba mokslinių straipsnių redaktore UAB „VTEX“.

Bibliography

- [1] D.R. Anderson. Green's function for a third-order generalized right focal problem. *J. Math. Anal. Appl.*, **288**(1):1–14, 2003. Doi:10.1016/S0022-247X(03)00132-X.
- [2] D.R. Anderson, T.O. Anderson and M.M. Kleber. Green's function and existence of solutions for a functional focal differential equation. *Electron. J. Differential Equations*, **2006**(12):1–14, 2006.
- [3] U.M. Ascher, R.D. Russell and R.M. Mattheij. *Numerical Solution of Boundary Value Problems for Ordinary Differential Equations*. SIAM, 1995.
- [4] N.S. Bahvalov, H.P. Židkov and G.M. Kobel'kov. *Čislenyje Metody*. Laboratoriya Bazovyh Znanij, 2003. (in Russian)
- [5] A.V. Bitsadze and A.A. Samarskii. Some elementary generalizations of linear elliptic boundary value problems. *Dokl. Akad. Nauk SSSR*, **185**:739–740, 1969. (in Russian)
- [6] J.R. Cannon. The solution of the heat equation subject to specification of energy. *Quart. Appl. Math.*, **21**(2):155–160, 1963.
- [7] R. Čiegis, A. Štikonas, O. Štikonienė and O. Suboč. Stationary problems with nonlocal boundary conditions. *Math. Model. Anal.*, **6**(2):178–191, 2001.
Doi:10.1080/13926292.2001.9637157.
- [8] R. Čiegis, A. Štikonas, O. Štikonienė and O. Suboč. A monotonic finite-difference scheme for a parabolic problem with nonlocal conditions. *Differ. Equ.*, **38**(7):1027–1037, 2002.
Doi:10.1023/A:1021167932414.
- [9] R. Čiegis and N. Tumanova. Numerical solution of parabolic problems with non-local boundary conditions. *Numer. Funct. Anal. Optim.*, **31**(12):1318–1329, 2010.
Doi:10.1080/01630563.2010.526734.
- [10] R. Čiuپaila, Ž. Jesevičiūtė and M. Sapagovas. On the eigenvalue problem for one-dimensional differential operator with nonlocal integral condition. *Nonlinear Anal. Model. Control*, **9**(2):109–116, 2004.
- [11] D.G. Duffy. *Green's Functions with Applications*. Chapman & Hall/CRC Press, 2001.
- [12] A.V. Gulin, N.I. Ionkin and V.A. Morozova. Difference schemes with nonlocal boundary conditions. *Comput. Methods Appl. Math.*, **1**(1):62–71, 2001.
- [13] A.V. Gulin and V.A. Morozova. On the stability of a nonlocal finite-difference boundary value problem. *Differ. Equ.*, **39**(7):962–967, 2003.
Doi:10.1023/B:DIEQ.0000009192.30909.13.
- [14] A.V. Gulin, V.A. Morozova and N.S. Udvovichenko. Stability criterion for a family of non-local difference schemes. *Differ. Equ.*, **46**(7):973–990, 2010.
Doi:10.1134/S0012266110070050.
- [15] A.V. Gulin, V.A. Morozova and N.S. Udvovichenko. Stability of a nonlocal difference problem with a complex parameter. *Differ. Equ.*, **47**(8):1116–1129, 2011.
Doi:10.1134/S0012266111080064.

- [16] V.A. Il'in. Necessary and sufficient properties of being a basis of a subsystem of eigenfunctions and adjoint functions for Keldysh bundle for ordinary differential operators. *Dokl. Akad. Nauk SSSR*, **227**(4):796–799, 1976. (in Russian)
- [17] V.A. Il'in and E.I. Moiseev. Nonlocal boundary value problem of the first kind for a Sturm–Liouville operator in its differential and finite difference aspects. *Differ. Equ.*, **23**(7):803–810, 1987.
- [18] V.A. Il'in and E.I. Moiseev. Nonlocal boundary value problem of the second kind for a Sturm–Liouville operator. *Differ. Equ.*, **23**(8):979–987, 1987.
- [19] G. Infante. Eigenvalues of some non-local boundary-value problems. *Proc. Edinb. Math. Soc.* (2), **46**:75–86, 2003. Doi:10.1017/S0013091501001079.
- [20] G. Infante. Eigenvalues and positive solutions of ODEs involving integral boundary conditions. *Discrete Contin. Dyn. Syst.*, **2005**:436–442, 2005.
- [21] G. Infante and J.R.L. Webb. Positive solutions of some nonlocal boundary value problems. *Abstr. Appl. Anal.*, **18**:1047–1060, 2003. Doi:10.1155/S1085337503301034.
- [22] G. Infante and J.R.L. Webb. Three-point boundary value problems with solutions that change sign. *J. Integral Equations Appl.*, **15**(1):37–57, 2003.
Doi:10.1216/jiea/1181074944.
- [23] N.I. Ionkin. Difference schemes for one nonclassical problem. *Mosc. Univ. Comput. Math. Cybern.*, **2**:20–32, 1977. (in Russian)
- [24] N.I. Ionkin. The solution of a certain boundary value problem of the theory of heat conduction with a nonclassical boundary condition. *Differ. Uravn.*, **13**(2):294–304, 1977. (in Russian)
- [25] N.I. Ionkin and V.A. Morozova. Two dimensional heat equation with nonlocal boundary conditions. *Differ. Uravn.*, **36**(7):884–888, 2000. (in Russian)
- [26] F. Ivanauskas, T. Meškauskas and M. Sapagovas. Stability of difference schemes for two-dimensional parabolic equations with non-local boundary conditions. *Appl. Math. Comput.*, **215**(7):2716–2732, 2009. Doi:10.1016/j.amc.2009.09.012.
- [27] L.I. Kamynin. A boundary value problem in the theory of the heat conduction with nonclassical boundary condition. *Zh. Vychisl. Mat. Mat. Fiz.*, **4**(6):1006–1024, 1964. (in Russian)
- [28] Q. Ma. Existence of positive solutions for the symmetry three-point boundary-value problem. *Electron. J. Differential Equations*, **2007**(154):1–8, 2007.
- [29] R.Y. Ma. Existence of positive solutions for superlinear semipositone m -point boundary value problems. *Proc. Edinb. Math. Soc.* (2), **46**:279–292, 2003.
Doi:10.1017/S0013091502000391.
- [30] R.Y. Ma and B. Thompson. Global behavior of positive solutions of nonlinear three-point boundary value problems. *Nonlinear Anal.*, **60**(4):685–701, 2005.
Doi:10.1016/j.na.2004.09.044.

- [31] M. Sapagovas, A. Štikonas and O. Štikonienė. Alternating Direction Method for the Poisson Equation with Variable Weight Coefficients in an Integral Condition. *Differ. Equ.*, **47**(8):1163–1174, 2011. ISSN 0012-2661. Doi:10.1134/S0012266111080118.
- [32] S. Pečiulytė and A. Štikonas. On positive eigenfunctions of Sturm–Liouville problem with nonlocal two-point boundary condition. *Math. Model. Anal.*, **12**(2):215–226, 2007. Doi:10.3846/1392-6292.2007.12.215-226.
- [33] S. Pečiulytė, O. Štikonienė and A. Štikonas. Sturm–Liouville problem for stationary differential operator with nonlocal integral boundary condition. *Math. Model. Anal.*, **10**(4):377–392, 2005. Doi:10.1080/13926292.2005.9637295.
- [34] A.A. Samarskii. *The Theory of Difference Schemes*. Marcel Dekker, Inc., New York, Basel, 2001.
- [35] A.A. Samarskij and A.V. Gulin. *Chislennye Metody*. Nauka, Moskva, 1989. (in Russian)
- [36] A.A. Samarskij and E.S. Nikolaev. *Metody Reshenija Setochnyh Uravnenij*. Nauka, Moskva, 1978. (in Russian)
- [37] M. Sapagovas. *Diferencialinių lygčių kraštiniai uždaviniai su nelokaliosiomis sąlygomis*. Mokslo aidai, Vilnius, 2007. (in Lithuanian)
- [38] M.P. Sapagovas. The eigenvalues of some problem with a nonlocal condition. *Differ. Equ.*, **38**(7):1020–1026, 2002. Doi:10.1023/A:1021115915575.
- [39] M. Sapagovas. On the stability of a finite-difference scheme for nonlocal parabolic boundary-value problems. *Lith. Math. J.*, **48**(3):339–356, 2008. Doi:10.1007/s10986-008-9017-5.
- [40] M. Sapagovas and R. Čiegis. The numerical solution of some nonlocal problems. *Lith. Math. J.*, **27**(2):348–356, 1987.
- [41] M. Sapagovas and R. Čiegis. On some boundary problems with nonlocal conditions. *Differ. Uravn.*, **23**(7):1268–1274, 1987. (in Russian)
- [42] M. Sapagovas, G. Kairytė, O. Štikonienė and A. Štikonas. Alternating direction method for a two-dimensional parabolic equation with a nonlocal boundary condition. *Math. Model. Anal.*, **12**(1):131–142, 2007. Doi:10.3846/1392-6292.2007.12.131-142.
- [43] A. Skučaitė, K. Skučaitė-Bingelė, S. Pečiulytė and A. Štikonas. Investigation of the spectrum for the Sturm–Liouville problem with one integral boundary condition. *Nonlinear Anal. Model. Control*, **15**(4):501–512, 2010.
- [44] I. Stakgold. *Green's Functions and Boundary Value Problems*. Wiley–Interscience Publications, New York, 1979.
- [45] A. Štikonas. Investigation of characteristic curve for Sturm–Liouville problem with nonlocal boundary conditions on torus. *Math. Model. Anal.*, **16**(1):1–22, 2011. Doi:10.3846/13926292.2011.552260.
- [46] Y. Sun. Eigenvalues and symmetric positive solutions for a three-point boundary-value problem. *Electron. J. Differential Equations*, **2005**(127):1–7, 2005.

- [47] L.X. Truong, L.T.P. Ngoc and N.T. Long. Positive solutions for an m -point boundary-value problem. *Electron. J. Differential Equations*, **2008**(111):1–11, 2008.
- [48] V.S. Vladimirov. *Equations of Mathematical Physics*. Nauka, Moskva, 1981. (in Russian)
- [49] J.R.L. Webb. Remarks on positive solutions of some three point boundary value problems. In *Proceedings of the Fourth International Conference on Dynamical Systems and Differential Equations, May 24–27, 2002*, pp. 905–915. Wilmington, NC, USA, 2003.
- [50] J.R.L. Webb. Positive solutions of some higher order nonlocal boundary value problems. *Electron. J. Qual. Theory Differ. Equ.*, **29**:1–15, 2009.
- [51] J.R.L. Webb and G. Infante. Nonlocal boundary value problems of arbitrary order. *J. London Math. Soc.*, **79**:238–258, 2009. Doi:10.1112/jlms/jdn066.
- [52] J.R.L. Webb, G. Infante and D. Franco. Positive solutions of nonlinear fourth order boundary value problems with local and nonlocal boundary conditions. *Proc. Roy. Soc. Edinburgh Sect. A*, **138**:427–446, 2008. Doi:10.1017/S0308210506001041.
- [53] D. Xie, Y. Liu and C. Bai. Green’s function and positive solutions of a singular n th-order three-point boundary value problem on time scales. *Electron. J. Differential Equations*, **2009**(38):1–14, 2009.
- [54] B. Yang. Positive solutions of a third-order three-point boundary-value problem. *Electron. J. Differential Equations*, **2008**(99):1–10, 2008.
- [55] Z. Zhao. Positive solutions for singular three-point boundary-value problems. *Electron. J. Differential Equations*, **2007**(156):1–8, 2007.
- [56] Z. Zhao. Exact solutions of a class of second-order nonlocal boundary value problems and applications. *Appl. Math. Comput.*, **215**:1926–1936, 2009.
Doi:10.1016/j.amc.2009.07.043.

Svetlana ROMAN

GRYNO FUNKCIJOS UŽDAVINIAMS
SU NELOKALIOSIOMIS KRAŠTINĖMIS SĄLYGOMIS

Daktaro disertacija

Fiziniai mokslai (P 000),
Matematika (01 P)

Svetlana ROMAN

GREEN'S FUNCTIONS FOR BOUNDARY-VALUE PROBLEMS
WITH NONLOCAL BOUNDARY CONDITIONS

Doctoral Dissertation

Physical sciences (P 000),
Mathematics (01 P)

2011 11 07. 2 sp. l. Tiražas 60 egz.

Išleido VšĮ Vilniaus universiteto leidykla
Universiteto g. 1, LT-01122 Vilnius.
Interneto svetainė: <http://www.leidykla.vu.lt>.